arXiv:1004.4519v2 [math-ph] 27 Apr 2010

Quantum conditional entropy for infinite-dimensional systems[*]

A. A. Kuznetsova
Moscow State University
kelit@list.ru

Abstract

In this paper a general definition of quantum conditional entropy for infinite-dimensional systems is given based on recent work of Holevo and Shirokov [3] devoted to quantum mutual and coherent informations in the infinite-dimensional case. The properties of the conditional entropy such as monotonicity, concavity and subadditivity are also generalized to the infinite-dimensional case.

1 The definition

In this paper a general definition of quantum conditional entropy for infinite-dimensional systems is given based on recent work of Holevo and Shirokov [3] devoted to quantum mutual and coherent informations in the infinite-dimensional case. The necessity of such a generalization is clear in particular from the study of Bosonic Gaussian channels and was stressed also in [7]. We refer to [3] for some notations and preliminaries. Let $A, \dots$ be quantum systems described by the corresponding Hilbert spaces $\mathcal{H}_A, \dots$, and let Φ be a channel from A to B with the Stinespring isometry $V : \mathcal{H}_A \longrightarrow \mathcal{H}_B \otimes \mathcal{H}_E$. Let ρ_{AR} be a purification of a state ρ_A with the reference system R and let ρ_{BRE} be the pure state obtained by action of the operator $V \otimes I_R$. The *mutual*

---

[*] Work partially supported by RFBR grant 09-01-00424.

information of the quantum channel Φ at the state ρ_A is defined similarly to finite-dimensional case (cf. [1], [2]) as

$$I(\rho_A, \Phi) = H(\rho_{BR} \| \rho_B \otimes \rho_R),$$

and *coherent information* at the state ρ_A *with finite entropy* $H(\rho_A)$ is defined as follows [3]

$$I_c(\rho_A, \Phi) = I(\rho_A, \Phi) - H(\rho_A).$$

It is then shown that the above-defined quantity satisfies the inequalities

$$-H(\rho_A) \leq I_c(\rho_A, \Phi) \leq H(\rho_A)$$

and the identity

$$I_c(\rho_A, \Phi) + I_c(\rho_A, \widetilde{\Phi}) = 0, \tag{1}$$

where $\widetilde{\Phi}$ is the complementary channel.

Let A, C be two (in general, infinite-dimensional) systems and ρ_{AC} a state such that $H(\rho_C) < \infty$. We define the *conditional entropy* as

$$H(C|A) = H(\rho_C) - H(\rho_{AC} \| \rho_A \otimes \rho_C), \tag{2}$$

where $H(\cdot\|\cdot)$ is the relative entropy which takes its values in $[0, +\infty]$, so that $H(C|A)$ is well defined as a quantity with values in $[-\infty, +\infty)$. If in addition $H(\rho_A) < \infty$, one recovers the standard formula $H(C|A) = H(AC) - H(A)$.

Now let B be infinite-dimensional system so that arbitrary state ρ_{BC} can be purified to a pure state ρ_{ABC}. Consider the channel $\Phi = \mathrm{Tr}_A$ from AB to B so that $\widetilde{\Phi} = \mathrm{Tr}_B$, then

$$I_c(\rho_{AB}, \mathrm{Tr}_B) \equiv I(\rho_{AB}, \mathrm{Tr}_B) - H(\rho_{AB}) = H(\rho_{AC} \| \rho_A \otimes \rho_C) - H(\rho_C) = -H(C|A). \tag{3}$$

Similarly

$$I_c(\rho_{AB}, \mathrm{Tr}_A) \equiv I(\rho_{AB}, \mathrm{Tr}_A) - H(\rho_{AB}) = H(\rho_{BC} \| \rho_B \otimes \rho_C) - H(\rho_C) = -H(C|B). \tag{4}$$

From the results of [3] concerning the coherent information, we then obtain that under the condition $H(\rho_C) < \infty$ both $H(C|A), H(C|B)$ are well defined and satisfy

$$|H(C|A)| \leq H(\rho_C), \quad |H(C|B)| \leq H(\rho_C), \tag{5}$$

$$H(C|A) + H(C|B) = 0, \tag{6}$$

2 Properties of conditional entropy

Proposition 1. *Let* ρ_{AB} *be such a state that* $H(\rho_A) < \infty$. *The function* $H(A|B)$ *has the following properties:*

1.monotonicity: *the inequality*

$$H(A|BC) \leq H(A|B). \tag{7}$$

holds for any systems A, B, C.

2. concavity in ρ_{AB} : *if* $\rho_{AB} = \alpha\rho^1_{AB} + (1-\alpha)\rho^2_{AB}, \alpha \in [0, 1]$, *then*

$$H(A|B) \geq \alpha H(A^1|B^1) + (1-\alpha)H(A^2|B^2). \tag{8}$$

3. subadditivity: *the inequality*

$$H(AB|CD) \leq H(A|C) + H(B|D). \tag{9}$$

holds for any systems A, B, C, D *such that* $H(\rho_A) < \infty$, $H(\rho_B) < \infty$.

Proof. 1. To prove monotonicity we rewrite the inequality (7) using the definition (2), i.e.

$$H(\rho_A) - H(\rho_{ABC}\|\rho_{BC} \otimes \rho_A) \leq H(\rho_A) - H(\rho_{AB}\|\rho_B \otimes \rho_A).$$

which is equivalent to

$$H(\rho_{AB}\|\rho_B \otimes \rho_A) \leq H(\rho_{ABC}\|\rho_{BC} \otimes \rho_A),$$

and the last inequality holds by monotonicity of the relative entropy with respect to taking the partial trace.

2. Let P^A_n, P^B_k be arbitrary increasing sequences of finite rank projectors in the spaces $\mathcal{H}_A, \mathcal{H}_B$, strongly converging to the operators I_A, I_B respectively. Consider the sequence of states

$$\rho^{nk}_{AB} = \lambda^{-1}_{nk}(P^A_n \otimes P^B_k \rho_{AB} P^A_n \otimes P^B_k), \lambda_{nk} = \mathrm{Tr}(P^A_n \otimes P^B_k)\rho_{AB},$$

with the partial states ρ^{nk}_A, ρ^{nk}_B.

Let us show first that

$$\lim_{n,k\to\infty} H(A_{nk}|B_{nk}) = H(A|B). \tag{10}$$

By the definition (2) $H(A_{nk}|B_{nk})=H(\rho_A^{nk})-H(\rho_{AB}^{nk}\|\rho_B^{nk}\otimes\rho_A^{nk})$. By using lower semi-continuity of the von Neumann entropy [4], we obtain

$$\lim_{n,k\to\infty}\inf H(\lambda_{nk}\rho_A^{nk})\geq H(\rho_A).$$

On the other hand, [5, lemma 4] implies

$$H(\lambda_{nk}\rho_A^{nk})=H(P_n^A(\mathrm{Tr}_B I\otimes P_k^B\rho_{AB}I\otimes P_k^B)P_n^A)\leq H(\mathrm{Tr}_B I\otimes P_k^B\rho_{AB}I\otimes P_k^B).$$

Further, by the dominated convergence theorem for entropy [4] we have

$$\lim_{n,k\to\infty}H(\mathrm{Tr}_B I\otimes P_k^B\rho_{AB}I\otimes P_k^B)=H(\rho_A),$$

since $\mathrm{Tr}_B(I\otimes P_k^B\rho_{AB}I\otimes P_k^B)\leq\rho_A$ and $H(\rho_A)<\infty$. Thus by the theorem about the limit of the intermediate sequence we obtain

$$\lim_{n,k\to\infty}H(\lambda_{nk}\rho_A^{nk})=H(\rho_A),\ \text{hence,}\ \lim_{n,k\to\infty}H(\rho_A^{nk})=H(\rho_A).$$

Then we prove that $\lim\limits_{n,k\to\infty}H(\rho_{AB}^{nk}\|\rho_A^{nk}\otimes\rho_B^{nk})=H(\rho_{AB}\|\rho_A\otimes\rho_B)$. Consider the following values

$$H_{nk}=H(\rho_{AB}^{nk}\|\rho_B^{nk}\otimes\rho_A^{nk})=H(\rho_A^{nk})+H(\rho_B^{nk})-H(\rho_{AB}^{nk}),$$

$$\widetilde{H}_{nk}=H\left(\rho_{AB}^{nk}\|\eta_k^{-1}P_k^B\rho_BP_k^B\otimes\mu_n^{-1}P_n^A\rho_AP_n^A\right)=-H(\rho_{AB}^{nk})-$$

$$-\mathrm{Tr}\left(\rho_A^{nk}\right)\log\left(\mu_n^{-1}P_n^A\rho_AP_n^A\right)-\mathrm{Tr}\left(\rho_B^{nk}\right)\log\left(\eta_k^{-1}P_k^B\rho_BP_k^B\right),$$

$$\text{where } \mu_n=\mathrm{Tr}P_n^A\rho_A,\ \eta_k=\mathrm{Tr}P_k^B\rho_B.$$

By using again [5, lemma 4] we have

$$\begin{aligned}\lim_{n,k\to\infty}\widetilde{H}_{nk} &= \lim_{n,k\to\infty}H\left(P_n^A\otimes P_k^B\rho_{AB}P_n^A\otimes P_k^B\|P_k^B\rho_BP_k^B\otimes P_n^A\rho_AP_n^A\right)\\ &= H(\rho_{AB}\|\rho_B\otimes\rho_A).\end{aligned}$$

We will prove that $\lim_{n,k\to\infty}H_{nk}=H(\rho_{AB}\|\rho_A\otimes\rho_B)$ by considering the difference $H_{nk}-\widetilde{H}_{nk}$. After some calculation we obtain that the difference tends to zero:

$$\lim_{n,k\to\infty}\left(H_{nk}-\widetilde{H}_{nk}\right)=\lim_{n,k\to\infty}\left(H\left(\rho_A^{nk}\|\mu_n^{-1}P_n^A\rho_AP_n^A\right)+H\left(\rho_B^{nk}\|\eta_k^{-1}P_k^B\rho_BP_k^B\right)\right)=0.$$

The last double limit is equal to zero since it follows from [5, lemma 4] that

$$0 \leq H\left(\lambda_{nk}\rho_A^{nk}\|P_n^A\rho_A P_n^A\right) = H\left(P_n^A \mathrm{Tr}_B(I_A\otimes P_k^B\rho_{AB}I_A\otimes P_k^B)P_n^A\|P_n^A\rho_A P_n^A\right) \leq$$

$$\leq H\left(\mathrm{Tr}_B(I_A\otimes P_k^B\rho_{AB}I_A\otimes P_k^B)\|\rho_A\right),$$

and [3, lemma 7] implies

$$\lim_{n,k\to\infty} H\left(\mathrm{Tr}_B(I_A\otimes P_k^B\rho_{AB}I_A\otimes P_k^B)\|\rho_A\right) = H(\rho_A\|\rho_A) = 0.$$

Thus, the theorem about the limit of the intermediate sequence implies

$$\lim_{n,k\to\infty} H\left(\lambda_{nk}\rho_A^{nk}\|P_n^A\rho_A P_n^A\right) = \lim_{n,k\to\infty} H\left(\rho_A^{nk}\|\mu_n^{-1}P_n^A\rho_A P_n^A\right) = 0.$$

Similarly we obtain that the second summand of the difference $H_{nk} - \widetilde{H}_{nk}$ also tends to zero:

$$\lim_{n,k\to\infty} H\left(\rho_B^{nk}\|\eta_k^{-1}P_k^B\rho_B P_k^B\right) = 0.$$

Finally, we have

$$\lim_{n,k\to\infty} H(\rho_{AB}^{nk}\|\rho_B^{nk}\otimes\rho_A^{nk}) = H(\rho_{AB}\|\rho_B\otimes\rho_A), \text{ hence } \lim_{n,k\to\infty} H(A_{nk}|B_{nk}) = H(A|B).$$

To prove the concavity of the function H(A—B) let us consider

$$\rho_{AB} = \alpha\rho_{AB}^1 + (1-\alpha)\rho_{AB}^2, \alpha\in[0,1].$$

Also consider again

$$\rho_{AB}^{nk} = \lambda_{nk}^{-1}P_n^A\otimes P_k^B\rho_{AB}P_n^A\otimes P_k^B =$$

$$= \frac{\alpha(P_n^A\otimes P_k^B\rho_{AB}^1 P_n^A\otimes P_k^B) + (1-\alpha)(P_n^A\otimes P_k^B\rho_{AB}^2 P_n^A\otimes P_k^B)}{\alpha\mathrm{Tr}(P_n^A\otimes P_k^B\rho_{AB}^1) + (1-\alpha)\mathrm{Tr}(P_n^A\otimes P_k^B\rho_{AB}^2)} = \frac{\theta_{nk}^1\rho_{AB}^{1nk} + \theta_{nk}^2\rho_{AB}^{2nk}}{\theta_{nk}^1+\theta_{nk}^2},$$

$$\theta_{nk}^1 = \alpha\mathrm{Tr}P_n^A\otimes P_k^B\rho_{AB}^1,\ \rho_{AB}^{1nk} = \frac{\alpha P_n^A\otimes P_k^B\rho_{AB}^1 P_n^A\otimes P_k^B}{\theta_{nk}^1},$$

$$\theta_{nk}^2 = (1-\alpha)\mathrm{Tr}P_n^A\otimes P_k^B\rho_{AB}^2,\ \rho_{AB}^{2nk} = \frac{(1-\alpha)P_n^A\otimes P_k^B\rho_{AB}^2 P_n^A\otimes P_k^B}{\theta_{nk}^2}.$$

Due to concavity of the condition information on the finite rank states we can write that

$$H(A_{nk}|B_{nk}) \geq \frac{\theta^1_{nk}}{\theta^1_{nk}+\theta^2_{nk}} H(A^1_{nk}|B^1_{nk}) + \frac{\theta^2_{nk}}{\theta^1_{nk}+\theta^2_{nk}} H(A^2_{nk}|B^2_{nk})$$

Taking the limit and using (10) for both parts of inequality we obtain the assertion of (8). Thus, concavity is proved.

3. A direct verification shows that in the finite-dimensional case

$$H(AB|CD) = H(A|CD) + H(B|CD) - (H(A|CD) - H(A|BCD)), \quad (11)$$

which implies

$$H(AB|CD) \leq H(A|CD) + H(B|CD), \quad (12)$$

since the value in brackets in (11) is nonnegative. We will prove the inequality (12) in infinite-dimensional case, the subadditivity property (9) can be derived from (12) by using monotonicity of the conditional entropy.

Let $\rho_{ABCD} = \rho \in \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C \otimes \mathcal{H}_D$ be a state with $H(\rho_A) < \infty, H(\rho_B) < \infty$. Consider an arbitrary increasing sequence of finite rank projectors $\{P^{CD}_l\}$ which strongly converges to the operator I_{CD}. Also consider the sequence of states

$$\rho^l = \tau_l^{-1}(I_A \otimes I_B \otimes P^{CD}_l)\rho(I_A \otimes I_B \otimes P^{CD}_l), \tau_l = \mathrm{Tr}(I_A \otimes I_B \otimes P^{CD}_l)\rho.$$

The relation (11) (which implies (12)) holds for ρ^l since all the summands in (11) are finite.

We will show that $H(A^l|CD^l) \to H(A|CD)$. Actually, using the definition (2) we have

$$H(A^l|CD^l) = H(\rho^l_A) - H(\rho^l_{ACD} \| \rho^l_{CD} \otimes \rho^l_A).$$

It follows from the dominated convergence theorem for entropy [4] that

$$\lim_{l\to\infty} H(\rho^l_A) = \lim_{l\to\infty} H\left(\tau_l^{-1}\mathrm{Tr}_{BCD}(I_A \otimes I_B \otimes P^{CD}_l)\rho\right) = H(\rho_A).$$

Further, consider the values

$$H_l = H(\rho^l_{ACD} \| \rho^l_{CD} \otimes \rho^l_A) = -H(\rho^l_{ACD}) + H(\rho^l_A) + H(\rho^l_{CD}).$$

and

$$\widetilde{H}_l = H(\rho^l_{ACD} \| \rho^l_{CD} \otimes \rho_A) = H\left(\tau_l^{-1} I_A \otimes P^{CD}_l[\rho_{ACD}] I_A \otimes P^{CD}_l \| \rho_A \otimes \tau_l^{-1} P^{CD}_l[\rho_{CD}] P^{CD}_l\right) =$$

$$= -H(\rho^l_{ACD}) + H(\rho^l_{CD}) + \mathrm{Tr}\left(\tau_l^{-1}\mathrm{Tr}_{CD} I_A \otimes P_l^{CD}[\rho_{ACD}]\right)(-\log\rho_A).$$

Then [5, lemma 4] implies that

$$\lim_{l\to\infty} \widetilde{H}_l = H(\rho_{ACD}\|\rho_A\otimes\rho_{CD}).$$

On the other hand, after the calculation and using [3, lemma 7] we obtain

$$\lim_{l\to\infty}(\widetilde{H}_l - H_l) = \lim_{l\to\infty} H\left(\tau_l^{-1}\mathrm{Tr}_{CD} I_A \otimes P_l^{CD}[\rho_{ACD}]\|\rho_A\right) = 0.$$

This implies that

$$\lim_{l\to\infty} H(\rho^l_{ACD}\|\rho^l_A\otimes\rho^l_{CD}) = H(\rho_{ACD}\|\rho_A\otimes\rho_{CD}),\ \text{hence,} \lim_{l\to\infty} H(A^l|CD^l) = H(A|CD).$$

In the similar way we obtain $H(B^l|CD^l) \to H(B|CD)$ and $H(AB^l|CD^l) \to H(AB|CD)$. Thus the statement (12) is proved in infinite-dimensional case, hence, the subadditivity property holds. □

The following observation is due to M. E. Shirokov.

Proposition 2. *Let $\mathcal{A}$ be a subset of $\mathfrak{S}(\mathcal{H}_{AC})$ such that the von Neumann entropy is continuous on the set $\mathcal{A}^C = \mathrm{Tr}_A\mathcal{A} \subset \mathfrak{S}(\mathcal{H}_C)$. Then the function $\rho_{AC} \mapsto H(C|A)$ is continuous on the set $\mathcal{A}$.*

Proof. Let $\{\rho_n\} \subset \mathcal{A}$ be a sequence converging to a state $\rho_0 \in \mathcal{A}$. By the well known results of purification theory there exists a corresponding sequence of purifications $\{\hat{\rho}_n\} \subset \mathfrak{S}(\mathcal{H}_{ABC})$ converging to a purification $\hat{\rho}_0 \in \mathfrak{S}(\mathcal{H}_{ABC})$ of the state ρ_0. The sequence $\{\mathrm{Tr}_C\hat{\rho}_n\}$ converges to $\mathrm{Tr}_C\hat{\rho}_0$ and $\lim_{n\to+\infty} H(\mathrm{Tr}_C\hat{\rho}_n) = H(\mathrm{Tr}_C\hat{\rho}_0)$ by the condition (since $H(\mathrm{Tr}_C\hat{\rho}_n) = H(\mathrm{Tr}_{AB}\hat{\rho}_n)$). Proposition 4 in [3] implies

$$\lim_{n\to+\infty} I_c(\mathrm{Tr}_C\hat{\rho}_n, \mathrm{Tr}_B) = I_c(\mathrm{Tr}_C\hat{\rho}_0, \mathrm{Tr}_B).$$

□

The author is grateful to A.S. Holevo and M.E. Shirokov for suggesting the problem and for discussions and the help in preparing the manuscript.

References

[1] *Adami C., Cerf N.J.* Capacity of noisy quantum channel // Phys.Rev. A. 1997. V.56 P.3470-3485; arXiv: quant-ph/9609024.

[2] *Barnum H., Nielsen M.A. Schumacher B.* Information transmission through a noisy quantum channel // Phys. Rev. A. 1998. V.57 P.4153-4175; arXiv:quant-ph/9702049.

[3] *Holevo A. S., Shirokov M. E.* Mutual and coherent informations for infinite-dimensional quantum channels. arXiv:1004.2495[quant-ph].

[4] *Wehrl A.* General properties of entropy// Reviews of Modern Physics. 1978. V.50, N.2, P. 221-260.

[5] *Lindblad G.* Expectations and Entropy Inequalities for Finite Quantum Systems// Commun. Math. Phys. 1974. V. 39. P. 111-119.

[6] *Holevo A. S., Shirokov M. E.* On approximation of infinite-dimensional quantum channels. Probl. Inform. Transmission. 2008. v. 44. n. 2. p. 3-22; arXiv:0711.2245[quant-ph].

[7] *Fabian Furrer, Johan Eberg, and Renato Renner* Min- and Max-Entropy in Infinite Dimensions //arXiv:1004.1386[quant-ph].